Phases of Matter Vocabulary

Word	Definition
Avogadro's Hypothesis	Equal volumes of two ideal gases under the same conditions of
	temperature and pressure will contain equal number of molecules.
Boiling	The transition of a liquid into a gas at the boiling point
Condensing	The transition of a gas into a liquid at the boiling point.
Deposition	The transition of a gas into a solid.
Equilibrium	The condition that exists when the rates of two opposing changes are equal.
Evaporating	The transition of the surface molecules of a liquid into a gas below the boiling point.
Freezing	The transition of a liquid into a solid at the freezing point.
Gas	A phase of matter characterized by the complete dissociation of matter particles from each other with the distances between the particles very large in comparison to the size of the particles and no attractive forces between them.
Heat of Fusion	The energy required to melt a gram of solid at its melting point.
Heat of Vaporization	The energy required to boil a gram of liquid at its boiling point.
Ideal Gas	A gas in which the molecules are infinitely small and far apart, the molecules travel with a straight-line motion, all collisions have no net loss of energy (elastic), there are no attractive forces between molecules and the speed of the molecules is directly proportional to the Kelvin temperature. Gases are most ideal at high temperature and low pressure.
Liquid	A phase of matter characterized by matter loosely organized yet kept together by intermolecular or ionic attractive forces.
Melting	The transition of a solid into a liquid at the melting point.
Pressure	Force exerted over an area.
Solid	A phase of matter characterized by matter arranged in regular geometric patterns called crystal lattices with only vibration motion, no relative motion.
Specific Heat	The energy required to heat one gram of a substance by one Kelvin.
Sublimation	The transition of a solid into a gas.
Vapor pressure	The pressure exerted by vapor in a vapor-liquid mixture in a closed system at equilibrium.
Vapor-Liquid Equilibrium	A system where the rate of evaporation equals the rate of condensing.

1) Phases and Phase Change (HW: p. 18, 19)


Essential Question: How does the proximity of atoms or molecules to each other affect the properties they exhibit?

Positive and negative charges attract each other. Attractive forces between molecules are called **intermolecular attractive forces**. The strength of these forces determines what phase of matter a substance is in at a given temperature. Substances with weak attractive forces (London dispersion) between their molecules tend to be gases at room temperature, and substances with strong attractive forces (ionic) tend to be solids at room temperature. Phases of matter are simply stages of attraction. Gases are made of molecules with no attractive forces, allowing the molecules to fly freely past each other. Liquids are made of molecules with stronger attractive forces, allowing the molecules to flow past each other, but still stay together. Solids are made of molecules or ions with strong attractive forces, which lock the molecules into a crystal lattice where the particles are free to vibrate, but they cannot move relative to each other.

PROPERTIES OF PHASES:


SOLIDS:

- a) Molecules, atoms or ions are arranged into a regular, geometric pattern called a crystal lattice.
- b) Molecules, atoms or ions vibrate in place. They do not move relative to each other.
- c) Definite shape, definite volume.




LIQUIDS:

- a) Molecules, atoms or ions can flow past each other.
- b) Because of intermolecular attractive forces (IMAF), there is resistance to flow, called viscosity.
- c) Viscosity increases as temperature decreases and IMAF strength increases.
- d) Liquid molecules near the surface can escape into the vapor phase below the boiling point, a process called evaporation.
- e) Shape of container (in absence of gravity, they form a perfect sphere), definite volume.




GASES:

- a) Gas molecules are extremely far apart compared to the size of the molecules.
- b) Gas molecules travel in a straight line until they collide with something.
- c) Collisions are elastic which means they don't lose any kinetic energy in the collision.
- d) Gas molecules move faster when it's hotter (higher Kelvin temperature)
- e) The gas phase is the only phase that is affected by changes in pressure.
- f) They spread out to take the shape and volume of whatever container they are put into.



Phase Change Diagram:




Phase Equilibrium: boiling and condensing both occur at the boiling point (100 °C for water), freezing and melting both occur at the melting point (0 °C for water). During the phase change, both phases exist at equilibrium.

EQUILIBRIUM: A condition where the rates of opposing changes are equal. So, a substance at the melting (freezing) point is melting at the same rate that it is freezing.

This is why water melts ABOVE 0° C and freezes BELOW) 0° C. AT 0° C, a mixture of water and ice will not change either way. If you have a sealed flask that contains 20.0 grams of ice and 40.0 grams of liquid water and keep it at 0° C, come back in a day, year, month, century, millennium, eon...there will still be 20.0 grams of ice and 40.0 grams of water. That's equilibrium!


Sublimation: attractive forces between molecules are so weak that heating a solid causes it to go directly into the gas phase. There are two common substances that undergo this change; $CO_2(s)$, also known as dry ice, and $I_2(s)$.

Water also undergoes sublimation! Place a fresh batch of ice cubes in the freezer and leave them there for a few months. You will notice that the ice cubes will have shrunk! You also might notice a rime of frost (ice crystals) on the walls of the freezer! Sometimes you have to scrape this frost out, or it builds up thick. What happened is that the surface molecules of water in the ice sublimed (turned into water vapor), hit the very cold walls of the freezer and froze there! When the gas turns directly back into a solid, as happens here (and on your car's windshield on very cold mornings), it is called **deposition**.




Phase Change Diagrams (Heating Curve):


When a solid (**A**) is heated, its temperature increases until the melting point is reached (**B**)(for H_2O : 0°C or 273 K). The solid then absorbs potential energy as it melts (BC). During the melting process, the temperature remains constant until all of the solid has melted (**C**). Once the substance is completely in the liquid phase, the temperature will increase as the liquid is continuously heated (**CD**). Once the boiling point has been reached (**D**) (for H_2O : $100^{\circ}C$ or 373 K), the temperature will remain constant as the substance turns from a liquid into a gas, a process called boiling (or vaporization)(**DE**). Once the substance is entirely in the gas phase (**E**), the temperature will increase as heat is continually applied (**EF**). The phase changes in this scenario are both **endothermic**, because heat is being constantly applied.



Heating Curve For Water

This graph shows what happens to the temperature of a sample of H₂O (s) as it is heated from 200 K to 420 K.





The melting point of water is 0 °C, which is 273 K. The boiling point of water is 100°C, which is 373 K.

How many minutes pass from the first appearance of the liquid phase until the substance is entirely in the gas phase?

The liquid first appears at **B** (4 minutes). It is entirely in the gas phase at **E** (34 minutes). 34-4 = **30 minutes**

How many minutes will it take for this substance to undergo melting?

The solid starts to melt at point **B** (4 minutes), and it is finished melting at point **C** (8 minutes). 8-4 = 4 minutes

For how many minutes is the water completely in a phase made of a crystal lattice?

The substance is a crystal lattice in the **solid** phase, from **A** (0 minutes) to **B** (4 minutes). 4-0 = **4 minutes**

What line segment represents when H₂O is both in the liquid AND the gas phase?

The substance is in both the liquid and the gas phase during **boiling**, which takes place from **D** (20 minutes) to **E** (34 minutes). 34 - 20 = 14 minutes

For how many minutes is the water completely in a phase with no definite shape or volume?

The **gas** phase has no definite shape or volume. The substance is completely in the gas phase from **E** (34 minutes) until **F** (36 minutes). 36-34 = 2 minutes

How many minutes will it take for the water to boil, once the boiling point temperature has been reached?

The substance reaches the boiling point at **D** (20 minutes). Boiling takes until **E** (34 minutes). 34-20 = **14 minutes**

2) Energy Required for Phase Change (HW: p. 20-21)

Essential Question: How does the defroster in your car work?

- 1) **Melting and boiling are endothermic** because energy has to be constantly added in order for the change to be completed.
- **2) Condensing and freezing are exothermic** because energy has to be constantly removed in order for the change to be completed.

HEAT OF FUSION: The amount of heat energy needed to melt one gram of solid at the substance's melting point. For water, the heat of fusion is 334 J/g. You can find this information on Reference Table B. In order to calculate the heat (in joules) needed to melt a sample of a substance, use the equation $\mathbf{q} = \mathbf{mH_f}$. You can find this equation on Reference Table T. If you find that you have frost on your windshield in the morning, turn on the defroster. It will heat the frost to the melting point, and then melt the ice by adding heat of fusion. The same thing is true of your microwave if you are using it to defrost frozen food. The microwave heats the food to the melting point and then adds heat of fusion to melt it so that you can use the food item to make yourself some dinner!

Heat of fusion can also be used for determining how many joules are required to freeze a liquid at the freezing point (which is the same temperature at the melting point). Use the same calculation. The only difference is that when you melt a substance, you have to ADD the heat of fusion to the solid, and when you freeze the substance, you have to REMOVE heat from the liquid. When you put water into a freezer, the freezer removes heat from the water to get it to the freezing point, then removes heat of fusion to turn it all into ice.

How many joules does it take to melt 100. grams of water at its melting point?

 $q = mH_f = 100$. grams X 334 J/gram = **33 400 Joules**

How many joules does it take to freeze 50.0 grams of water at its freezing point?

 $q = mH_f = 50.0 \text{ grams X } 334 \text{ J/gram} = 16 700 \text{ Joules}$

HEAT OF VAPORIZATION: the amount of heat energy needed to boil one gram of liquid at the substance's boiling point. For water, the heat of vaporization is 2260 J/g. You can find this information on Reference Table B. In order to calculate the heat (in joules) needed to boil a sample of a substance, use the equation $\mathbf{q} = \mathbf{mH_{V}}$. You can find this equation on Reference Table T. If you have a solution and need to recover the solute dissolved in it, put the solution in an evaporating dish and heat it up over a Bunsen burner or hot plate. Once the boiling point of water has been reached, heat of vaporization boils off all of the water, so you are left with pure solute.

Heat of vaporization can also be used for determining how many joules are released when a gas condenses back into a liquid. Use the same calculation. Which causes greater injury, having boiling water at 100°C splashed on you, or steam at 100°C? The water is hot and your skin absorbs that heat...but the steam is also hot, and as it hits your skin, it releases heat of vaporization as well, causing greater injury. Ask to see my steam burn scar sometime.

Heat of vaporization is absorbed from your skin by sweat as it evaporates, which removes heat from your skin, leaving you feeling cooler. It is our own personal, if slightly gross, air conditioning system.

How many joules are required to boil 100. grams of water at its boiling point?

 $q = mH_V = 100$. grams X 2260 J/gram = **226 000 Joules**


How many joules does it take to condense 50.0 grams of water at its boiling point?

 $q = mH_V = 50.0 \text{ grams } X 2260 \text{ J/gram} = 113 000 \text{ Joules}$

Determining Heat Required to Heat A Substance Through Phase Changes

Problem: How many joules are required to heat 100.0 grams of H₂O (s) from -20.00°C to 130.00°C?

Solution: Water undergoes two phase changes between those two temperatures.



During segments AB, CD and EF, temperature is changing. Therefore, the equation $q = mc\Delta T$ can be used.

The specific heat of liquid water is 4.18 J/g°C. The specific heat of solid and gaseous water is 2.09 J/g°C.

During segment BC, water is melting. Heat of fusion is required here (334 J/g).

During segment DE, water is boiling. Heat of vaporization is required here (2260 J/g).

So, this calculation will be done in five steps!

1) -20.00° C to 0.00° C (AB): $q = mc\Delta T = 100.0 \text{ g X } 2.09 \text{ J/g}^{\circ}$ C X 20.0° C = **4180 J** to heat it to melting point

2) melting (BC): $q = m H_{f=} 100.0 g X 334 J/g = 33400 J to melt it$

3) 0.00° C to 100.00° C (CD): $q = mc\Delta T = 100.0 \text{ g X } 4.18 \text{ J/g}^{\circ}$ C X 100.00° C = **41800 J** to heat it to boiling point

4) boiling (DE): $q = m H_v = 100.0 g X 2260 J/g = 226000 J to boil it$

5) 100.00° C to 130.00° C (EF): $q = mc\Delta T = 100.0 \text{ g X } 2.09 \text{ J/g}^{\circ}$ C X 30.00° C = **6270 J to heat it to 130°C**

To find the total amount of energy needed to perform this change, simply add up all the joules!

TOTAL AMOUNT OF ENERGY NEEDED = 311650 J, which rounds to 312000 J

Problem: If the energy is added at the rate of 2000 J/ min, how long will this take?

312000 J / 2000 J/min = **156 min**

3) Gases and Pressure (HW: p. 22-24)

Essential Question: What assumptions do we have to make when we set a standard for physical properties?

KINETIC-MOLECULAR THEORY (IDEAL GAS BEHAVIOR)

- 1) Gases are made molecules that are extremely tiny and far apart from each other. This is why gases can be compressed.
- 2) Gas particles move in a constant straight-line motion.
- **3) Any collisions a gas particle makes will be elastic that is, there no loss of energy**. Each time a gas molecule collides against an obstacle, it bounces off with the same speed that it hit with.
- **4) There are no intermolecular attractive force between gas particles**. All substances have attractive forces, but the distances between the molecules are so great, they don't come into play. Gases with London dispersion forces behave the most like an ideal gas would, and gases with hydrogen bond attractions behave the least like an ideal gas would.
- **5)** The average speed of the particles is directly proportional to the kelvin (absolute) temperature. In other words, the gas molecules move faster when it's hotter.

IDEAL GAS BEHAVIOR: Under conditions of HIGH temperature and LOW pressure, molecules behave more like an "ideal" gas. The smaller the gas molecule is, the more ideal it will behave, therefore, hydrogen and helium are the gases that are the most ideal.

AVOGADRO'S HYPOTHESIS: equal numbers of particles of a gas will occupy equal volume under the same conditions of temperature and pressure. This is because gas molecules will all spread out to an equal degree at the same temperature and pressure. This means that any math we want to do with gases can be applied to all gases equally!

All three gases are at 298 K and 101.3 kPa

CO₂

NH₃

^{1 L} О₂

They all contain equal numbers of molecules.

DEVIATIONS FROM THE IDEAL GAS LAW - WHERE AND WHY?

This law works well under **standard conditions** of temperature and pressure (STP). (Ref. Table A).

Standard temperature is 0°C, or 273 K.

Standard pressure is 1.000 atm or 101.3 kPa or 760.0 mmHg (torr).

PRESSURE: force exerted over an area. The gases in the atmosphere exert pressure because of Earth's gravity. The pressure exerted is 14.7 pounds per square inch (psi). This means that over every square inch of surface (including YOU), on average, at sea level, the air molecules in our atmosphere are pushing with 14.7 pounds of force. You are rather lucky in that your internal pressure (and the pressure in your cells) is also about14.7 pounds per square inch. You can feel this pressure if you drive up a mountain or ride in an airplane. As you go up in the atmosphere, there is less air above you exerting pressure. To equalize to this lower pressure, your ears pop!

Pressure is measured in atmospheres (atm), kilopascals (kPa) or millimeters of mercury (mmHg).

CONVERSIONS: 14.7 psi = 1.000 atm = 101.3 kPa = 760.0 mmHg

Convert 2.35 atm to kPa:

(2.35 atm) X (101.3 kPa/-atm) = 238 kPa

Convert 123.4 kPa to atm:

(123.4 kPa) / (101.3 kPa / atm) = 1.218 atm

<u>VAPOR PRESSURE</u>: the pressure exerted by a liquid's vapor in a sealed container at vapor-liquid equilibrium at a given temperature. The vapor pressure of a liquid is not dependent on the mass or volume of the liquid. The vapor pressure of a liquid can be found on **Reference Table H**. **The stronger the attractive forces are between liquid molecules**, **the lower the vapor pressure is**.

Substances that have high vapor pressures evaporate very quickly. Gasoline is a mixture of liquids with very high vapor pressures, you can even see the gas evaporate as you pump it into your gas tank. Alcohol and acetone (nail polish remover) also evaporate very quickly. Water evaporates fairly slowly, because the hydrogen bonding keeps the liquid molecules more attracted to each other than in the other liquids, meaning the vapor molecules can't escape as easily. Substances that evaporate readily are called **volatile**. Most of these liquids are also quite flammable. The vapor burns even faster than the liquid does. This is why you are not supposed to get back into your car when you are pumping gas until you are done...rubbing your feet on the carpet in your car can cause you to build up a static charge, and if you touch your car, you might cause a spark that could cause an explosion.

BOILING POINT: the temperature at which a liquid's vapor pressure equals the pressure exerted on the liquid by outside forces. Use Reference Table H to determine a liquid's boiling point. Boiling point increases as exerted pressure is increased.

NORMAL BOILING POINT: the boiling point of a liquid under a pressure of 1.000 atmosphere. Substances with higher boiling points have stronger attractive forces holding molecules to each other in the liquid phase, making it require more energy to overcome those attractions and permit boiling.

- 1) The normal boiling point of water at sea level is 100° C. At this elevation, the atmosphere is exerting 1.000 atm of pressure.
- 2) If one takes a ride into the mountains, the amount of air above you decreases, and so does the pressure. Since there is less pressure to keep water in the liquid state, its boiling point decreases. This is why there are special cooking directions for higher elevations. You can't hard-boil an egg when the boiling point is too low!
- 3) Let's say you have some dried beans and you want to cook them up. To do so, soak them in water for a while, and then add heat. It takes 90 minutes to cook pinto beans (the ones used for refried beans) on the stove at normal atmospheric pressure. Put them in a pressure cooker instead, where the lid is sealed shut so the pressure can just rise, rise and rise), and the boiling point of the water will increase so much that now it will only take 5 to 7 minutes to cook those same beans!
- 4) On the surface of Mars, the average atmospheric pressure is 0.6 to 1 kPa, depending on where you are. At this pressure, the boiling point of water and the melting point of water are the same! If you were to bring an ice cube on to Mars and heat it, it would undergo sublimation and never form a liquid! That's why any liquid water on Mars must be under the ground, where there is higher pressure to support the liquid phase.

How To Use Table H:

A) What is the vapor pressure of ____ at ___ °C?

What is the vapor pressure of ethanol at 40°C? Start at the 40°C point on the X axis, trace a line up to the ethanol curve, then shoot across and read the vapor pressure off of the Y axis. The vapor pressure is 17 kPa.

B) What is the boiling point of _____ at a pressure of ____ kPa?


What is the boiling point of water at a pressure of 30 kPa? Start at the 30 kPa point on the Y axis, trace a line to the water curve, then shoot down and read the temperature off the X axis. The boiling point is 70°C.

C) What is the normal boiling point of ____?

The normal boiling point is the temperature at which a liquid boils under standard pressure, 101.3 kPa. There is a dashed line going across the table at 101.3 and even labeled "101.3 kPa". Follow the 101.3 kPa line to the curve of your interest (in this case, water), and shoot down to get the boiling point from the X axis. As expected, water has a normal boiling point of 100°C!

Notice that the scales for the X and Y axis are different. Each box on the Y axis is 10 kPa, but every box on the X axis is 5°C.

Table H Vapor Pressure of Four Liquids



4) Partial Pressure and Effusion (HW: p. 25, 26)

Essential Question: Why can you smell dog poop on the corner of your lawn sooner after the dog does the deed in the summer than in the winter?

DALTON'S LAW OF PARTIAL PRESSURES: The total pressure exerted by a mixture of gases is equal to the sum of the pressures exerted by each of the gases in the mixture.

Examples:

1) What is the total pressure of a mixture of O_2 (g), N_2 (g) and NH_3 (g) if the pressure of the O_2 (g) is 20. kPa, N_2 (g) is 60. kPa and the NH_3 (g) is 15 kPa?

$$P_{\text{total}} = P_{\text{O2}} + P_{\text{N2}} + P_{\text{NH3}} = 20$$
. kPa + 60. kPa + 15 kPa = **95 kPa**

2) A mixture of 1 mole of O_2 and 2 moles of N_2 exerts a pressure of 150. kPa. What is the partial pressure of each gas?

There are 3 moles of gas total. 150. kPa / 3 moles = 50.0 kPa/mole. Since there is 1 mole of O_2 its partial pressure is 50.0 kPa. Since there are 2 moles of O_2 its partial pressure is O_2 its par

3) A mixture of 30.0% He and 70.0% Ar exerts a pressure of 150 kPa at 25°C. What is the partial pressure of each gas?

For **He**: 150. kPa X 0.300 (the decimal form of 30.%) = **45.0 kPa**

For **Ar**: 150. kPa X 0.700 (the decimal form of 70.%) = **105 kPa**

4) A sample of NH_3 (g) is decomposed into its component elements. If the pressure of the nitrogen gas produced equals 40.0 kPa, what would the pressure of the hydrogen gas?

Since the formula of ammonia shows a 1:3 ratio of nitrogen to hydrogen, the pressure of the hydrogen produced will be three times greater than the pressure of the nitrogen.

40.0 kPa X 3 = **120. kPa**

Effusion: as molecular mass (and density) decreases and the temperature increases, gases spread out faster. If a light molecule and a heavy molecule are put in a race at the same temperature, the lighter gas would be able to move faster and the heavier molecule would move slower. Think of trying to push two vehicles parked in neutral on a level surface. One is a Geo Metro, the other is a Mack eighteen-wheeler semi. Now, get behind each one in turn and push. Which one goes faster?

A closed container of a mixture of chlorine, fluorine, neon and helium gases is opened so the gases can escape. Place the gases in order of increasing rate of effusion. Using Reference Table S, we find the densities of the gases to be $CI = 0.003214 \text{ g/cm}^3$, $F = 0.001696 \text{ g/cm}^3$, $Ne = 0.0009 \text{ g/cm}^3$ and $He = 0.000179 \text{ g/cm}^3$. Therefore, putting them in order of increasing rate of effusion, we have **He, Ne, F and CI**.

5) The Gas Laws (HW: p. 27-30)

Essential Question: How can models of particle behavior be used to make predictions?

The Gas Laws are relationships between temperature, pressure and volume of a gas. Gas law equations are used to determine what effect changing one of those variables will have on any of the others.

What are the units for pressure, volume and temperature?

Pressure: atm or kPa Volume: mL or L Temperature: K

The gas laws are based on one fundamental truth called Avagadro's Hypothesis:

EQUAL VOLUMES OF TWO SAMPLES OF IDEAL GASES CONTAIN EQUAL NUMBERS OF PARTICLES UNDER THE SAME CONDITIONS OF TEMPERATURE AND PRESSURE.

Consider two 4.00 L containers, each at 298 K and 1.00 atm. Container A holds nitrogen gas, Container B holds carbon dioxide gas. If container A holds 2.00 moles of nitrogen gas, how many moles of carbon dioxide must be present in container B?

Since both containers contain equal volumes of gases under the same conditions of temperature and pressure, Avagadro's Hypothesis holds that both containers will contain equal number of molecules, and therefore equal number of molecules. **Therefore, if container A holds 2.00 moles of gas, so must container B**.

This means that all gases behave more or less equally to changes in temperature, pressure and volume, so one equation can be used to describe these changes and applied equally to all gases, if one assumes that they exhibit ideal behavior.

Do equal volumes of gases under the same conditions of temperature and pressure have the same MASS? Why or why not?

No, because each element has it's own unique atomic mass.

To solve gas law problems, follow these easy steps!

1) Get rid of the words! Create a little data table and look through the numbers. If the units are mL or L, it's volume! Pick out V_1 and V_2 . If the units are atm or kPa, it's pressure! Pick out P_1 and P_2 . If the units are K, it's temperature! Pick out P_1 and P_2 . You may not have to pick out all of the variables, as sometimes one of them is constant and you won't have to worry about it.

HINT: If you see the unit °C, that is temperature, too. In gas laws, temperature MUST be in Kelvin, so simply add 273 and get your T1 and T2 in Kelvin!


- 2) Write the gas law equation down. Actually write it right there on the page. In the blank spot. Go ahead...
- **3) Circle the variable you are trying to solve for, and use basic algebra to rearrange the equation**. Don't be afraid to ask for help.
- **4)** Substitute the numbers into your newly rearranged equation, and enter them into your calculator. Don't forget to push the "=" or "Enter" button!
- **5) Round your answer off.** Since this is all multiplication and division, round your answer off to the same number of sig figs as the data you put in that had the fewest number of sig figs! Don't forget to put your unit after your answer!

RELATIONSHIPS BETWEEN VARIABLES OF PRESSURE, VOLUME AND TEMPERATURE

1) Pressure vs. Volume (Constant Temperature): as pressure is increased, volume is decreased (inverse relationship)

BOYLE'S LAW (pressure vs. volume at constant temperature)

$$P \times V = K$$
 $P_1V_1 = K$, $P_2V_2 = K$, therefore $P_1V_1 = P_2V_2$



A 10.0 L (V_1) sample of gas is trapped in a cylinder with a movable piston at constant temperature. The pressure of the gas in this cylinder is 1.0 atm (P_1). The cylinder is compressed until the pressure doubles to 2.0 atm (P_2). The new volume of the gas will be 5.0 L (V_2). If pressure is doubled, then volume is cut in half.

A sample of gas occupies a volume of 2.00 L at STP. If the pressure is increased to 2.00 atm at constant temperature, what is the new volume of the gas?

Set up a table: (since P₁ is given as STP, look up its value on Table A...it's 1.00 atm!)

$$P_1 = 1.00 \text{ atm}$$
 $V_1 = 2.00 \text{ L}$ $P_2 = 2.00 \text{ atm}$ $V_2 = x$

Then rearrange the Combined Gas Law Equation to solve for the desired variable, omitting the variable that is held constant:


 $P_1V_1 = P_2V_2$, solve for V_2 gives a rearranged equation of $P_1V_1/P_2 = V_2$

Then plug and chug! [(1.00 atm)(2.00 L)] / (2.00 atm) = 1.00 L

2) Volume vs. Temperature (Constant Pressure): as temperature is increased, volume is increased (direct relationship)

CHARLES' LAW (volume vs. temperature at constant pressure)

$$V/T = K$$
 $V_1/T_1 = K$, $V_2/T_2 = K$, therefore $V_1/T_1 = V_2/T_2$



A 1.0 L (V_1) sample of gas is trapped in a cylinder with a moveable piston at constant pressure. The temperature of the gas in this cylinder is 200. K (T_1) . The cylinder is heated until the temperature doubles to 400. K (T_2) . The new volume of the gas will be 2.0 L (V_2) . If temperature is doubled, then volume is doubled, too.

The temperature MUST be in Kelvin. If it is given in °C, you must add 273 to the temperature before solving the problem.

A sample of gas occupies a volume of 5.00 L at 300. K. If the temperature is doubled under constant pressure, what will the new volume of the gas be?

Set up a table:

$$V_1 = 5.00 L$$

$$I_1 = 300. K$$

$$V_1 = 5.00 L$$
 $T_1 = 300. K$ $V_2 = x$ $T_2 = 600. K(double T_1)$


Then rearrange the Combined Gas Law Equation to solve for the desired variable, omitting the variable that is held constant:

$$V_1$$
 V_2 = , solve for V_2 gives a rearranged equation of $V_1T_2/T_1 = V_2$ T_1 T_2

Then plug and chug! [(5.00 L)(600. K)]/(300. K) = 10.0 L

<u>3) Temperature vs. Pressure (Constant Volume):</u> as temperature is increased, pressure is increased (direct relationship)

GAY-LUSSAC'S LAW (pressure vs. temperature at constant volume) P/T = K P_1/T_1 = K, P_2/T_2 = K, therefore P_1/T_1 = P_2/T_2



A sample of gas is trapped in a rigid cylinder of fixed volume at 300. K (T_1) . The pressure of the gas in this cylinder is 2.0 atm (P_1) . The cylinder is heated until the temperature doubles to 600. K (T_2) . The new pressure of the gas will be 4.0 atm (P_2) . If temperature is doubled, then the pressure is doubled too.

The temperature MUST be in Kelvin. If it is given in °C, you must add 273 to the temperature before solving the problem.

A 10.0 L sample of gas in a rigid container at 1.00 atm and 200. K is heated to 800. K. Assuming that the volume remains constant, what is the new pressure of the gas?

Set up a table:

$$P_1 = 1.00 \text{ atm}$$
 $T_1 = 200. \text{ K}$

$$P_2 = x$$

$$T_2$$
 = 800. K

Then rearrange the Combined Gas Law Equation to solve for the desired variable, omitting the variable that is held constant:

$$P_1$$
 P_2 = , solve for P_2 gives a rearranged equation of $P_1T_2/T_1 = P_2$ T_1 T_2

Then plug and chug! [(1.00 atm)(800. K)]/(200. K) = 4.00 atm

COMBINED GAS LAW

These three gas laws can be combined to form the **Combined Gas Law**, as found on Reference Table T:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$P = \text{pressure}$$

$$V = \text{volume}$$

$$T = \text{tempera}$$

$$P = \text{pressure}$$

 $T = \text{temperature}(\mathbf{K})$

If a variable is constant, it may be ignored. Set up a table for the values of the variables and solve for the unknown. There might not BE anything held constant. In that case, use all of the variables!

A 2.00 L sample of gas at STP is heated to 500. K and compressed to 200. kPa. What is the new volume of the gas?

Set up a table:

$$P_1 = 101.3 \text{ kPa } V_1 = 2.00 \text{ L}$$
 $T_1 = 273. \text{ K}$ $P_2 = 200. \text{ kPa } V_2 = x$ $T_2 = 500. \text{ K}$

$$T_1 = 273$$
. K

$$P_2 = 200$$
. kPa $V_2 = 2$

$$T_2 = 500. K$$

Then rearrange the Combined Gas Law Equation to solve for the desired variable:

$$P_1V_1$$
 P_2V_2 = , solve for V_2 gives a rearranged equation of $(P_1V_1T_2)$ / (P_2T_1) = V_2 T_1 T_2

Then plug and chug! [(101.3 kPa)(2.00 L)(500. K)] / [(200. kPa)(273 K) = 1.86 L

A 2.00 L sample of gas at 1.00 atm and 300. K is heated to 500.K and compressed to a volume of 1.00 L. What is the new pressure of the gas?

Set up a table:

$$P_1 = 1.00 \text{ atm}$$
 $V_1 = 2.00 \text{ L}$ $T_1 = 300. \text{ K}$ $P_2 = x$ $V_2 = 1.00 \text{ L}$ $T_2 = 500. \text{ K}$

$$T_1 = 300. K$$

$$P_2 = x$$

$$V_2 = 1.00 I$$

$$T_0 = 500 \text{ K}$$

Then rearrange the Combined Gas Law Equation to solve for the desired variable:

$$P_1V_1$$
 P_2V_2 = , solve for P_2 gives a rearranged equation of $(P_1V_1T_2)$ / (V_2T_1) T_1 T_2

Then plug and chug! [(1.00 atm)(2.00 L)(500. K)] / [(1.00 L)(300. K)] = 3.33 atm

A 2.00 L sample of gas at 300. K and a pressure of 80.0 kPa is placed into a 1.00 L container at a pressure of 240. kPa. What is the new temperature of the gas?

Set up a table:

$$P_1 = 80.0 \text{ kPa}$$
 $V_1 = 2.00 \text{ L}$ $T_1 = 300. \text{ K}$ $P_2 = 240. \text{ kPa}$ $V_2 = 1.00 \text{ L}$ $T_2 = x$

$$T_1 = 300. k$$

$$P_2 = 240$$
. kPa $V_2 = 1.00$ L

$$T_2 = \lambda$$

Then rearrange the Combined Gas Law Equation to solve for the desired variable:

$$P_1V_1$$
 P_2V_2 = , solve for T_2 gives a rearranged equation of $(P_2V_2T_1)$ / (P_1V_1) T_1 T_2

Then plug and chug! [(240. kPa)(1.00 L)(300. K)] / [(80.0 kPa)(2.00 L)] = 450. K

IDEAL GAS LAW

The pressure and volume of a gas are proportional to the number of moles of gas and the Kelvin temperature. The equation can be derived as follows:

From the Combined Gas Law, we have

PV/T = K, K = nR (n is number of moles, R is a proportionality constant)

Since one mole of gas exerts a pressure of 1.00 atm and occupies a volume of 22.4 L at 273 K, R (the proportionality constant) can be derived as follows:

(1 atm)(22.4 L)/(1 mole)(273 K) = R

R = 0.0821 atm-L/mol-K

This yields the IDEAL GAS LAW, which can be used to determine the pressure, volume, temperature or number of moles of gas if all of the other conditions are known, and none of the conditions have changed.

PV = nRT

P= Pressure (atm) V = Volume (L) n = moles R = 0.0821 atm-L/mol-K T = Temperature (K)

What is the pressure exerted by 3.00 moles of gas at a temperature of 300. K in a 4.00 L contaner?

PV = nRT, therefore P = nRT / V = [(3.00 moles)(0.0821 atm-L/mol-K)(300. K)] / (4.00L) =**18.5 atm**

What is the volume of a sample of gas if 5.00 moles if it exerts a pressure of 0.500 atm at 200. K?

PV = nRT, therefore V = nRT / P = [(5.00 moles)(0.0821 atm-L/mol-K)(200.K)] / (0.500 atm) =**164 L**

A sample of gas is contained in a cylinder with a volume of 10.0 L. At what temperature will 2.50 moles of contained gas exert 20.0 atm of pressure on the container?

PV = nRT, therefore T = PV / nR = [(20.0 atm)(10.0 L)] / [(2.50 moles)(0.0821 atm-L/mol-K)] = 974 K

A sample of gas contained in a cylinder of 5.00 L exerts a pressure of 3.00 atm at 300. K. How many moles of gas are trapped in the cylinder?

PV = nRT, so n = PV / RT = [(3.00 atm)(5.00 L)] / [(0.0821 atm-L/mol-K)(300. K)] =**0.609 moles**

If you are asked to determine the value of R using units other than atm, L, mol or K, then simply convert the different unit to its equivalent.

Determine the value of R in atm-mL/mol-K

First, convert molar volume (22.4 L/mol) to mL/mol by multiplying by 1000. mL/L.

R = (1 atm)(22400 mL) / (1 mol)(273 K) = 81.5 atm-mL/mol-K

Student Name:	Grades:	Phases,	Heat	Gases	DL & GL	Gas Laws
1) PHASES AND PHASE CHANGE HOMEWO)RK					

A) Multiple Choice Questions: Place your answer in the space in front of each question.

1) Which sample has molecules that flow past each other, yet are still attracted to each other? a) $C_6H_{12}O_6$ (s) b) $C_6H_{12}O_6$ (I) c) $C_6H_{12}O_6$ (g) 2) NaCl (I) is said to be boiling when it undergoes a phase change to a) solid b) liquid c) gas __3) What phase of matter has definite volume, but not definite shape? a) solid b) liquid c) gas 4) Motor oil viscosity is tested at different temperatures to see how effective it is at providing engine lubrication. As the temperature increases, what happens to the viscosity of the oil? a) increases b) decreases c) remains the same Explain, in terms of molecular motion and average kinetic energy:

Explain WHY this change is exothermic:

Explain WHY the other changes are NOT exothermic:

____5) What phase of matter has the strongest attractive forces?
a) solid
b) liquid
c) gas

Explain why attractive forces between particles of a substance determine what phase a substance is in at a given temperature:

B) A beaker that contains 100. g of substance X in the solid phase is heated constantly over a hot plate at the rate of 1000. J/min, generating the following data:

Time (min)	Temp (K)						
0	275	6	291	12	351	18	381
1	279	7	291	13	366	19	381
2	283	8	291	14	381	20	381
3	287	9	306	15	381	21	381
4	291	10	321	16	381	22	390
5	291	11	336	17	381	23	399

- 1) Draw a graph of the data above, placing the temperature on the Y-Axis, and time on the X-axis. Label the following information on the graph: where the three phases occur and what and where the phase changes are.
- 2) Since you are adding 1000. joules each minute, based on your graph (show your work):
- a) How many joules are required to melt the substance?
- b) How many joules are required to boil the substance?
- 3) How many minutes pass from the first moment the liquid phase forms until the substance is completely in the gas phase?
- 4) For how many minutes is the substance completely in the liquid phase?_____
- 5) For how many minutes does the substance exist completely as a crystal lattice?_____
- 6) For how many minutes is the substance in a phase that has no definite mass or volume?
- 7) While the substance is being heated in the liquid phase, what happens to the viscosity of the liquid?
- 8) Water melts at 273 K and boils at 373 K. Which substance, X or water, has stronger intermolecular attractive forces? Explain briefly.
- 9) Specific heat is the number of joules required to heat one gram of substance by one degree Celsius.
- a) Rearrange the Calorimetry equation to solve for C (specific heat):
- b) Given the information in the introduction, calculate the specific heat of Substance X when it is in the liquid phase.

2) Energy Required for Phase Change Homework

A) Multiple Choice Questions: Place your answer in the space in front of each question.

1) Which of the following phase changes requires heat of fusion to accomplish?

 $\overline{a) H_2O}$ (s) $\rightarrow H_2O$ (g)

b) $H_2O(g) \rightarrow H_2O(l)$

c) $H_2O(I) \rightarrow H_2O(g)$

d) $H_2O(s) \rightarrow H_2O(l)$

2) Which of the following phase changes is endothermic?

 $\overline{a) H_2O}$ (s) $\rightarrow H_2O$ (I)

b) $H_2O(g) \rightarrow H_2O(l)$

c) $H_2O(I) \rightarrow H_2O(s)$

d) $H_2O(g) \rightarrow H_2O(s)$

3) A mixture of 50.0 g of ice (H₂O (s)) and 30.0 g of water (H₂O (l)) sits in a sealed flask at 0°C.

_a) What will happen to the amount of ice in the flask if the mixture is left alone at 0° C?

a) increase

b) decrease

c) remain the same

Explain your answer:

____b) What will happen to the amount of ice in the flask if the temperature of the flask is lowered to -10°C?

a) increase

b) What will happen to the amount of ice in the flask if the temperature of the flask is lowered to -10°C?

c) remain the same


Explain your answer:

B) Calculate the number of joules required to (show correct numerical setup):

- a) melt 20.0 g of H_2O (s) at $0^{\circ}C$
- b) boil 30.0 g of H_2O (I) at $100^{\circ}C$
- c) freeze 200.0 g of H₂O (I) at 0°C
- d) boil 50.0 g of H_2O (g) at $100^{\circ}C$

C) Ice cubes are placed in a saucepan and heated at a constant rate over a stove.

- 1) Calculate the number of joules required to bring 512 g of water from -12.0° C to 115.0°C. (The specific heat of water in the solid and gas phase is 2.09 J/g·°C)
- a) Sketch a phase change diagram for the phase changes that occur between -12.0 °C and 115.0°C. Label the temperatures at which the phase changes occur. Then label each line segment with a letter (A, B, C, D, E, etc.).



- b) How many steps are needed to solve this problem?_____
- c) Use as spaces as you need below to solve this problem step by step. Label what line segment you are solving in the space to the left of the space provided to show your work:

Step 1:_____:

Step 2:_____:

Step 3:_____:

Step 4: :

Step 5:_____:

- d) Add up the energies required for each step and record your answer, rounded properly:
- e) If heat is added at the rate of 3000. J/min, how many minutes will it take for the entire process to occur?

3) Gases and Pressure Homework

A) Multiple Choice Questions: Place your answer in the space in front of each question.

a) real gas molecule b) real gas molecule c) real gas molecule	vay do real gases develos are extremely far apples have attractive force es move faster at higher travel in a straight li	es er temperatures	ehavior?	
2) Under wh i a) 100 K and 1 atm c) 300 K and 1 atm	ich conditions will O ₂	behave most ideally b) 200 K and 1 d) 400 K and 1	atm	
${\text{a) O}_2}$ 3) Which gas	s behaves most like a	an ideal gas at STP? c) H ₂	d) F ₂	
What quality of an i	deal gas does your cho	pice best match?		
a) at 100°C	b) below		s Peak (14110 feet above sea level) c) above 100°C	, water will boil:
5) Which sub	ostance on Reference	e Table H has the stro	ngest attractive forces holding its r	molecules
a) propanone	b) ethanol	c) water	d) ethanoic acid	
6) Which sar 100. kPa? a) 1.0 L sample of N b) 2.0 L sample of N c) 2.0 L sample of F		the same number of 0. kPa 0. kPa 0. kPa 0. kPa	needed to break molecules apart from molecules as a 2.0 L sample of O_2 (s	
Evoloin in terms of	Avagadra's Hypothesi	ie:		

Explain, in terms of *Avogadro's Hypothesis*:

1) 2.0 atm =		=		_kPa	
2) 1950. kPa =		=	·	_atm	
C) Why is the atmospheric pre- Explain, in terms of the <i>quantit</i>				it is in the bottom of De	ath Valley?
D) In which location will water Death Valley? Explain, using y					oottom of
E) Based on Reference Table H	I, What is the va	por pressure of	<u>:</u>		
a) ethanol @ 80°C?		_ kPa			
b) propanone @ 20°C?		_ kPa			
c) ethanoic acid @ 110°C?		_ kPa			
F) Based on Reference Table H	l, what is the boi	iling point of:			
a) ethanol under a pressure of 70). kPa? _		°C		
b) water under a pressure of 10.	kPa? _		°C		
c) ethanoic acid under a pressure	e of 120. kPa? _		°C		
G) Based on Reference Table I	l, what is the no	rmal boiling po	int of propan	one?	°C
H) Why do cooking directions elevations?	require you to co	ook your food l	onger at high	er elevations than at low	<u>'er</u>
I) A flask containing nothing of the pressure inside the flask as					nappen to

B) Perform the following conversions, show your work in the space provided:

J) EXTRA CREDIT!!! You have discovered two Earth-sized planets circling around the star Sigma Cygnii. Preliminary investigations through spectral analysis show that H₂O exists on both planets. Here are some other facts about the two planets:

Planet	Distance from Sigma Cygnii	Average Temperature Range	Average Atmospheric Pressure
Garellia	4.5 X 10 ⁸ km	-40°C – 20°C	50 kPa
Mongar	7.7 X 10 ⁷ km	60°C – 105°C	20 kPa

On which of these two planets does water exist in the liquid phase for the greatest temperature range? Explain, using information from Reference Table H to support your hypothesis.

4) Partial Pressure and Effusion Homework (all work must be shown for credit)

A) Dalton's Law Of Partial Pressures

1) Three gases are mixed into the same container. The partial pressures of the gases when mixed are 1.50 atm, 500. kPa and 300. kPa respectively. What is the total pressure of the gas mixture?
2) A mixture of gas A and gas B exerts a pressure of 400. kPa. If the partial pressure of gas B is 310. kPa, what is the partial pressure of gas A?
3) The earth's atmosphere consists of 78.0% N_2 , 20.0% O_2 , 1.0% Ar and 1.0% other trace gases. Assuming that the composition of the atmosphere is homogeneous, calculate the partial pressures of N_2 , O_2 and Ar under a total pressure of 101.3 kPa.
4) A mixture of one mole of gas A and two moles of gas B exerts a pressure of 99 kPa. What is the partial pressure of each gas?
5) When H ₂ O is decomposed, the resulting oxygen and hydrogen gas are trapped in a 5.0 L container. If the pressure of the oxygen is 300. kPa, what is the pressure of the hydrogen gas?
6) Hydrogen gas is trapped by water displacement in a test tube at 25°C. The total pressure of the gas in the test tube is 102.0 kPa. What pressure is exerted by the trapped hydrogen gas?

B) Graham's Law Of Effusion

b) Granam s Law Or Linus	1011	
1) Will the smell of baking Explain in terms of <i>molect</i>		our nose faster if the house you are in is at 20°C or 30°C?
		ole 10 meters in front of you. Perfume A has a density of 1.2 g/L h perfume will you smell first? Explain in terms of <i>molecular</i>
3) List the following in ord	ler of increasing rate	e (from slowest to fastest) of effusion:
DENSITY	MATERIAL	
 4.3 g/L 2.8 g/L 6.7 g/L	A B C	
4) a) List the following in c krypton and nitrogen.	order of increasing r	rate of effusion using Reference Table S: oxygen, xenon,
	these gases in the c	order that you did. Identify the reference table you got the
separate them on the basi strong attractive forces, w down on their journey thro particles and move faster the column and out to the	s of their intermoled which attracts other in ough the column. M through the column detector and emerg lose to being an idea	ixture of gases through a thin column of hollow tubing to cular attractive forces. There are particles in the column with molecules with strong attractive forces to it and slows them lolecules with weaker attractive forces are less attracted to these a. Helium is used as a "carrier" gas to push the other gases into les from the column before any of the other gases. Unlike al gas, helium is a noble gas, and will not chemically react with
a) Why, in terms of <i>molect</i> gases?	ular mass, does the	helium travel through the column faster than any of the other
b) Rocauso bolium goos #	arough the column t	izetor than any gae, how doos the etropath of its intermolecular
attractive forces compare		faster than any gas, how does the strength of its intermolecular gases in the mixture?

5) The Gas Laws Homework

(all work must be shown for credit)

1) A sample of hydrogen gas has a volume of 1.00 L at a pressure of 100. kPa. If the temperature is kept constant and the pressure is raised to 140. kPa, what is the new volume of the gas?
2) A gas sample occupies 10.0 mL at 1.00 atm of pressure. If the gas is put into a sealed 20.0 mL container and the temperature remains constant, what is the resulting pressure of the gas?
3) When 500. mL of hydrogen gas is heated from 30.0°C to 60.0°C at constant pressure, the volume of the gas at 60.0°C will change to:
4) One sunny day in January, you notice that your front left tire looks a bit low. You inflate it to the recommended 35.0 psi (241 kPa). The temperature that day is a nice 40.0°F (4.4°C). Late that night, you have to run an errand, but you notice the tire doesn't look as full as before. You look at the thermometer, which reads a brutal 12.0°F (-11.1°C). What is the pressure in your tire, assuming volume is constant?

Sig gas 29.	Natural gas pipelines are air pressure tested after construction and prior to being placed in service. Inificant changes in temperature can occur over the duration of the test. A 6" diameter, 30,000 foot long is main is tested at 13.6 atm for 24 hours. A thermometer placed near the main at the start of the test reads 4°C. At the end of the 24 hours period the pressure gauge reads 12.6 atm and the temperature is 21.1°C. erefore, the pressure dropped by 1.00 atm. That drop might have been due entirely to a leak, or due to the apperature dropor perhaps a little bit of both!
a)	How much of that pressure drop was due to the temperature drop?
b)	Is there a leak?
c)	If so, what is the pressure of the leak?
	A sample of gas at 200.K occupies a volume of 10.0 L at a pressure of 2.00 atm. To what temperature must gas be raised to double both the volume and the pressure?
7) A	A 2.00 L sample of oxygen exerts a pressure of 2.00 atm at 273 K. If the temperature is raised to 546 K and volume decreased to 1.00 L, what will be the final pressure on the gas?
8) A	A gas occupies 10.0 L of volume at STP. How much volume will it occupy at 2.00 atm and 300. K?

9) A sample of $\rm H_2$ gas is trapped by water displacement. The gas occupies a volume of 10.0 mL at a temperature of 273 K and a pressure of 1.00 atm. A low-pressure system passes by, lowering the pressure to 0.850 atm and dropping the temperature to 260. K. What will the new volume of gas be?
10) One mole of an ideal gas occupies 22.4 L of volume at STP. What would the molar volume be of an ideal gas at 80.0 kPa and 290. K?
11) A sample of gas has a volume of 20.0 L at STP. If the density of this gas is 1.40 g/L, what is the mass of this gas at a temperature of 20.0°C and 130.0. kPa?
12) A CO ₂ fire extinguisher contains 15.5 L of CO ₂ (g) under a pressure of 500. atm at 22.0°C. When fired, the pressure of the CO ₂ drops to 95.0 kPa, and the volume of CO ₂ released is 458 L. a) What is the temperature of the expelled CO ₂ ?
b) CO ₂ sublimes (undergoes sublimation) at a temperature of -78.5°C. What phase does the CO ₂ exit as?

Ideal Gas Law Problems

13) What volume will 21.6 moles of neon gas occupy under a pressure of 150. kPa and a temperature of 275 K?
14) A 50.0 L container is designed to withstand a maximum pressure of 225 atmospheres. If the container is holding 112 moles of nitrogen gas, what is the maximum temperature that the container can withstand before bursting?
15) How many moles of helium gas will occupy a volume of 52.0 L at STP?
16) Calculate R (the Universal Gas Constant) using the units torr, mL, mol, K, knowing that 1 mole of ideal gas occupies a volume of 22.4 L at STP.
17) One mole of oxygen gas has a mass of 32.0 grams. What pressure will 100. grams of oxygen exert in a 22.0 L container at 300. K?

Review Questions:

1) At what tempera a) At the freezing point, 27 c) At the boiling point, 273				ist?	
2) At what tempera a) At the freezing point, 27 c) At the boiling point, 273	ture will 3 K	the equilibrium H ₂ b) At the freez	O (s) ⇔ H₂O (I) exi ing point, 373 K	st?	
3) Which phase characters a) $CO_2(I) \rightarrow CO_2(g)$ b)			$_{2}$ (s) \rightarrow CO ₂ (I) d) $CO_2(s) \rightarrow CO_2(s)$	(g)
$\frac{\text{4) Which substance}}{\text{a) } C_6H_{12}O_6\text{ (s)} \qquad \text{b)}$			ged in a regular, g $I_{12}O_6$ (g) d		n called a crystal lattice?
${a) C_6 H_{12} O_6 (s)}$ 5) Which sample has been been been by) C ₆ H ₁₂ O ₆ (aq)	
6) What phase of many solid	natter has	s the strongest att b) liquid		veen the molecul) gas	les?
7) Turning a solid i a) melting, which is endoth c) freezing, which is endotl			ch is exothermic nich is exothermic		
8) How many joules		uired to melt 10.0 3340 J	grams of water at (c) 2260 J		d) 22 600 J
9) How many joule a) 334 J		quired to boil 10.0 3340 J	grams of water at o		d) 22 600 J
10) In what way do a) real gas molecules are v c) real gas molecules have11) Which of the fo	very smal attractive	ll. b) real e forces. d) real	gas molecules are gas molecules mov	ve faster as it gets	s hotter.
	Gas	Temp (K)	Pressure (atm)	Volume (L)	
	CO ₂	300	2	4	
	O ₂	250	1	3	
	CH ₄	300 250	1	7	
	1 12	200	1	<i>I</i>	

Gas	lemp (K)	Pressure (atm)	volume (L)
CO ₂	300	2	4
O_2	250	1	3
CH ₄	300	2	4
H_2	250	1	7

a) CO₂ and O₂

b) CO₂ and CH₄

c) O₂ and H₂

d) O₂ and CH₄

_12) A gas will behave most ideally under what conditions?

a) high T, low P

b) high T, high P

c) low T, high P

d) low T, low P

_13) Which of the following gases behaves most ideally at STP? b) CO_2 c) H_2

d) NH₃

Activity: Joules needed to melt a gram of ice

Materials: Styrofoam calorimeter cup, thermometer, 600 mL beaker, 100 mL graduated cylinder

Procedure:

- 1) Measure out exactly 100.0 mL of hot water in the graduated cylinder.
- 2) Pour the water into the calorimeter.
- 3) Measure and record the temperature.
- 4) Immediately place 2 ice cubes into the water and stir. As the ice melts, replace it.
- 5) Continue adding ice until the temperature does not drop any more, and no more ice melts. Record the final temperature.
- 6) Pour the water in the calorimeter into the 600 mL beaker until only the ice remains, using the thermometer as a dam to make sure that no ice gets into the beaker. Discard the ice.
- 7) Pour 100.0 mL of the water into the graduated cylinder and discard it (this represents the 100.0 mL of water you initially put into the calorimeter).
- 8) Record the volume of the remaining water (this represents the water that resulted from the melting of the ice) using the same graduated cylinder.

Data:					
Initial Temperature of Water in Calorimeter	°C				
Final Temperature of Water in Calorimeter	°C				
Volume of ice that melted	mL				
Calculations: The density of water at 25.0°C is 1.00 g/mL. 1) Determine the mass of the water that you put in the calorimeter. (density of water = 1.00 g/mL)					
2) Determine the mass of ice that melted. (density of v	water = 1.00 g/mL)				
3) Determine the temperature change of the water. (Δ	Γ = final temperature – initial temperature)				
1) Use calculations 1 and 3 to determine the number (q = m c Δ T)	of joules transferred from the water in the cup to the ice				

5) Determine the heat of fusion of ice in joules per gram of ice. (Answer to step 4 / answer to step 2)
Conclusions (in complete sentences or show all work, including units and sig figs): 1) The accepted value of the heat of fusion of ice is 334 joules per gram. Determine the percent error of your experiment. Show all work, units and proper rounding.
2) What are some possible sources of error for this experiment? How could they be lessened?
3) Describe the heat transfer between the ice and the water in terms of kinetic and potential energy change.